

Durham Research Online

Deposited in DRO:

04 April 2019

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Williams, Samual T. and Collinson, Wendy and Patterson-Abrolat, Claire and Marneweck, David G. and Swanepoel, Lourens H. (2019) 'Using road patrol data to identify factors associated with carnivore roadkill counts.', PeerJ., 7 . p. 6650.

Further information on publisher's website:

<https://doi.org/10.7717/peerj.6650>

Publisher's copyright statement:

© 2019 Williams et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, reproduction and adaptation in any medium and for any purpose provided that it is properly attributed. For attribution, the original author(s), title, publication source (PeerJ) and either DOI or URL of the article must be cited.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Using road patrol data to identify factors associated with carnivore roadkill counts

Samual T. Williams^{1,2,3}, Wendy Collinson⁴, Claire Patterson-Abrolat⁴, David G. Marneweck^{4,5} and Lourens H. Swanepoel¹

¹ Department of Zoology, University of Venda, Thohoyandou, South Africa

² Department of Anthropology, Durham University, Durham, United Kingdom

³ Institute for Globally Distributed Open Research and Education (IGDORE), Hoedspruit, South Africa

⁴ Endangered Wildlife Trust, Johannesburg, South Africa

⁵ Eugène Marais Chair of Wildlife Management, Mammal Research Institute, University of Pretoria, South Africa

ABSTRACT

As the global road network expands, roads pose an emerging threat to wildlife populations. One way in which roads can affect wildlife is wildlife-vehicle collisions, which can be a significant cause of mortality through roadkill. In order to successfully mitigate these problems, it is vital to understand the factors that can explain the distribution of roadkill. Collecting the data required to enable this can be expensive and time consuming, but there is significant potential in partnering with organisations that conduct existing road patrols to obtain the necessary data. We assessed the feasibility of using roadkill data collected daily between 2014 and 2017 by road patrol staff from a private road agency on a 410 km length of the N3 road in South Africa. We modelled the relationship between a set of environmental and anthropogenic variables on the number of roadkill carcasses, using serval (*Leptailurus serval*) as a model species. We recorded 5.24 serval roadkill carcasses/100 km/year. The number of carcasses was related to season, the amount of wetland, and NDVI, but was not related to any of the anthropogenic variables we included. This suggests that roadkill patterns may differ greatly depending on the ecology of species of interest, but targeting mitigation measures where roads pass through wetlands may help to reduce serval roadkill. Partnering with road agencies for data collection offers powerful opportunities to identify factors related to roadkill distribution and reduce the threats posed by roads to wildlife.

Subjects Conservation Biology, Ecology, Zoology

Keywords Road ecology, Human-wildlife conflict, Wildlife management, Wildlife-vehicle collision

INTRODUCTION

Roads, particularly in developing countries like South Africa, are integral to a country's development and prosperity (Karani, 2008; Keshkamati, Looijen & Zuidgeest, 2009). However, traffic can also have a direct negative impact on both people and wildlife (Verster & Fourie, 2018), with many species at risk from wildlife-vehicle collisions (WVCs), often resulting in an animal's death, or 'roadkill'. For the last three decades the field of road ecology has highlighted the negative impacts that roads and their associated users have on

Submitted 7 January 2019
Accepted 15 February 2019
Published 29 March 2019

Corresponding authors
Samual T. Williams,
samual.t.williams@gmail.com
Lourens H. Swanepoel,
lourens.swanepoel.univen@gmail.com

Academic editor
Stuart Pimm

Additional Information and
Declarations can be found on
page 11

DOI 10.7717/peerj.6650

© Copyright
2019 Williams et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

biodiversity, and their potential to affect wildlife populations (*Kioko et al., 2015; Laurance et al., 2015; Parchizadeh et al., 2018*).

Infrastructure development impacts biodiversity and ecosystems through exposing ecological habitats to disturbance and fragmentation (*Benítez-López, Alkemade & Verweij, 2010*). Land use, land cover, and connectivity within the landscape may change due to expanding road networks (*Perz et al., 2013; Liang et al., 2014*). Thus a detailed understanding of the factors involved in WVCs (*Gunson & Teixeira, 2015*) is required to implement successful mitigation strategies. Several studies in the United States of America (USA) and Europe have successfully quantified some of these factors (*Gunson & Teixeira, 2015; Rytwinski et al., 2015*). As a result, effective mitigation measures have been applied in these areas; habitat connectivity and accessibility are promoted, safe passage for animals using roads is facilitated because natural movements are encouraged and WVCs are reduced (*Jackson & Griffin, 2000; Forman et al., 2003; Grilo, Bissonette & Santos-Reis, 2009; Goosem, Izumi & Turton, 2001; Bager & Rosa, 2010*).

However, in developing countries, efforts to reduce wildlife mortality around main roads are often hampered due to a lack of research, with other priorities usually dictated by the country's socio-economic situation (*Collinson et al., 2015a*). For example, the collection of roadkill data by dedicated research teams can be extremely costly due to the high sampling effort required (*Abra et al., 2018*). This often limits the number of roadkill studies, and to date only a handful of studies have focussed on roadkill in Africa (*Collinson et al., 2015b*), and yet these data are vital in order to implement effective mitigation strategies (*Gunson & Teixeira, 2015*). This is unfortunate because while Africa is incredibly rich in biodiversity (*Mittermeier et al., 2011*), it also has the fastest growing human population (and associated infrastructure) in the world (*United Nations, 2015*), which could have serious environmental impacts such as habitat loss, degradation, and fragmentation (*Tilman et al., 2017*).

A citizen science approach, whereby scientific data are collected by members of the public is one way in which costly sampling strategies can be overcome (*Conrad & Hilchey, 2011*). For example, as a consequence of citizen science surveys used to monitor the status of birds in the United Kingdom (UK), the UK government introduced targets to reverse population declines identified by the surveys (*Gregory & Van Strien, 2010; Greenwood, 2012*). While citizen science can be a powerful tool for the collection of scientific data, biases in survey effort can hinder studies of variables such as roadkill rates. An alternative but underutilised source of data on roadkill rates that has the potential to offer more consistent and measurable survey effort, is road patrols. Many highway agencies conduct regular patrols in order to resolve issues that could affect road users. Establishing partnerships between researchers and road patrol agencies could therefore offer significant potential for more effective data collection (*Périquet et al., 2018*).

In this study, we used data collected by road patrol staff of the N3 Toll Concession (N3TC) to explore the potential for partnering with road agencies to conduct roadkill studies. In 2011, with the aim of managing the impact of roads on wildlife, a partnership was established between the Endangered Wildlife Trust (EWT) and the N3TC, an organisation that operates a 415 km of the N3 highway in South Africa. Following training from the

EWT, N3TC patrol staff began collecting data on roadkill incidents. An examination of total roadkill counts showed that the most common carnivore carcass found on the N3 was serval (*Leptailurus serval*). Since this is also one of the more threatened species in the dataset (Thiel, 2015), serval was selected as a suitable indicator species for the study.

The serval is a medium-sized carnivore weighing 8–12 kg (Skinner & Chimimba, 2005). It displays a preference for wetland habitats (Ramesh & Downs, 2015b), the degradation and loss of which constitute the principal threat to serval populations (Thiel, 2011). Although rodents make up a large proportion of serval diets, they are also known to feed on other small prey such as birds, reptiles, and insects (Ramesh & Downs, 2015a; Ramesh & Downs, 2015b). Despite declining numbers throughout their range (Ramesh & Downs, 2013), servals are listed as Least Concern on the IUCN Red List of Threatened Species (Thiel, 2015) and Near Threatened on the South African Red List (Ramesh et al., 2016). Home ranges are estimated to be between 8 and 38 km² (Geertsema, 1985; Ramesh, Kalle & Downs, 2015), and population densities can vary from approximately 6–100 individuals per 100 km² (Ramesh & Downs, 2013; Looock et al., 2018). Servals display largely crepuscular and nocturnal activity patterns (Thiel, 2011; Ramesh & Downs, 2013), but they can also be active during the day (Geertsema, 1985). Although roadkill is thought to present a serious threat to servals (Ramesh et al., 2016), there have been few attempts to quantify these threats or identify associated risk factors.

We used data collected by N3TC road patrol staff as part of routine road patrols to identify which factors were related to serval roadkill counts on the N3. Using a modelling approach, we evaluated the relationships between a set of environmental and anthropogenic variables. Our aim was to determine which variables, if any, were related to serval roadkill counts, and could be used to develop informed management strategies to help mitigate serval roadkill. This approach allowed us to test several hypotheses that could explain serval roadkill patterns (Table 1).

MATERIALS & METHODS

Study site

The study focussed on 410 km of the N3 Toll Route (hereafter referred to as the N3). The N3 is of strategic importance as it links Johannesburg, the country's largest city, with the port of Durban, and is a major route for the transport of goods between the two cities (Fig. 1). The N3 is classified as a national route, the highest category in the South African road network. Most segments of the national route network are maintained by the South African National Roads Agency (SANRAL), but portions are maintained by provincial, local, or private road authorities. The N3 has been managed by N3TC since 1999 (N3 Toll Concession, 2018). Mean annual traffic volume for a 415 km length of the N3 that encompasses the 410 km stretch studied was approximately 85 million vehicles between 2014 and 2017, and the road is generally two lanes wide in each direction, with speed limits of either 100 or 120 km/h. The N3 passes through three of the eleven South African biomes (Savanna, Grassland and Indian Ocean Coastal Belt (Mucina & Rutherford, 2006)), including urban landscapes, communal land and agricultural areas.

Table 1 Hypotheses relating to the relationships between serval roadkill counts and predictor variables included in the models.

Type	Variable name	Variable description	Direction of association	Rationale
Environmental	Season	Wet or dry season	More in dry	Serval may range over larger distances when water is scarce, making them more vulnerable to WVCs
	Wetland	Proportion of 10 km buffer composed of wetland	+	Serval would be more abundant in areas rich in their preferred habitat
	NDVI	Normalized difference vegetation index	+	Areas with a greater NDVI will have greater primary productivity, supporting greater densities of rodents and serval
	Guineafowl	Count of guineafowl carcasses	+	Guineafowl may be preyed upon by serval, increasing serval density
Anthropogenic	Traffic	Average number of vehicles per year	+	More vehicles present more opportunities for WVCs
	Speed	Average speed limit (km/h)	+	Faster cars will make collisions more difficult to avoid
	Road width	Average width of road (m)	+	Wider roads take longer to cross
	Infrastructure	Total number of infrastructure points such as bridges and underpasses	–	Bridges and underpasses may provide more opportunities for serval to cross roads safely




Figure 1 Map showing the location of the section of the N3 studied, serval roadkill carcass locations recorded from 2014 to 2017, and wetland within 10 km of the road. Note that wetland is shown as gaps in the black non-wetland areas within the 10 km buffer. The inset map on the left shows the study area in relation to serval range in South Africa (adapted from [Thiel \(2015\)](#)), and the inset map on the right (location shown in blue) shows a closer view of serval roadkill carcass locations in relation to wetlands.

Full-size [DOI: 10.7717/peerj.6650/fig-1](https://doi.org/10.7717/peerj.6650/fig-1)




Figure 2 Total seasonal serval roadkill counts collected along a 410 km section of the N3 in South Africa between 2014 and 2017.

[Full-size](#) DOI: [10.7717/peerj.6650/fig-2](https://doi.org/10.7717/peerj.6650/fig-2)

Data collection

N3TC road patrol staff collected field data every day between 01/01/2014 and 31/12/2017 as part of their routine patrol of a 410 km stretch of the N3 between Johannesburg and Durban (Fig. 2).

The road patrol staff involved in the study undertook annual training with the EWT in wildlife identification and the collection of roadkill data. The day-long training sessions also included training on the importance of conservation, an introduction to road ecology, and how to mitigate the effects of roads on wildlife, in addition to how to follow the data collection protocol. Participants each received a manual and a field guide to identifying animals killed on the road. Staff members from the EWT also accompanied N3TC staff on a sample of their patrols to provide additional training in the field, and to gain a deeper understanding of how the patrols work. Total contact time between the EWT and N3TC staff for training was approximately 14 days per year.

Patrols were conducted four times per day (twice in each direction) at a speed of approximately 100 km/h. Patrol teams mostly consisted of two observers, each allocated six short sections to ensure the entire road was covered effectively. Observers recorded the species of roadkill carcasses encountered, the date and time of the observation, and the carcass location using the nearest route marker, located at 200 m intervals along the road. Carcasses were removed from the road to avoid recounts (Collinson et al., 2014; Guinard, Prodon & Barbraud, 2015). We focussed on analysing records of serval and guineafowl (family Numididae), as these were among the most numerous roadkill carcasses. We classified roadkill observations collected between the 1st of October to the 31st of March as occurring during the wet season, and roadkill collected from the 1st of April to the 31st of September as occurring during the dry season (Cook, Reason & Hewitson, 2004).

We divided the N3 into 41 sampling units, each with a length of 10 km, which corresponded to the home range size of serval ([Geertsema, 1985](#); [Bowland, 1990](#); [Ramesh, Kalle & Downs, 2016](#)). We estimated habitat variables by first creating a buffer with a radius of 10 km around each sampling unit, and we used land cover data ([Geoterraimage, 2015](#)) to calculate the most common habitat type and the proportion of wetland within the buffer. We also obtained data from N3TC on traffic volumes, speed limits, and the locations of infrastructure such as bridges and underpasses. We measured road width at 410 randomly-generated locations (10 locations per sampling unit) using Copernicus Sentinel satellite imagery ([European Space Agency, 2017](#)), measured in QGIS v3.0 ([QGIS Development Team, 2018](#)). We used the same locations to sample normalized difference vegetation index (NDVI) provided at a resolution of 250 m every 16 days throughout the study period ([Didan, 2015](#)). Road shapefiles were obtained from OpenStreetMap ([OpenStreetMap contributors, 2017](#)).

Data analysis

We conducted all analyses in R v3.5.0 ([R Development Core Team, 2018](#)), and all code and data are publicly available ([Williams et al., 2018](#)). Within each sampling unit we summed the total number of serval carcasses, which we used as the response variable in our models. All other environmental and anthropogenic variables were also summarised for each sampling unit and used as potential predictors in the models ([Table 1](#)).

We modelled the relationship between serval roadkill counts and predictor variables by fitting a generalised linear mixed-effect model using the lme4 package ([Bates et al., 2015](#)). Exploratory analyses showed that negative binomial or Poisson distributions were the most plausible for our dataset. The negative binomial distribution with log link outperformed other model structures based on the Akaike Information Criterion (AIC) and visual inspection of a serval roadkill count histogram comparing distributions ([Fig. S1](#)), which was expected as the data were over-dispersed with an excess of zeros. This model structure was therefore used for the remaining analyses, following [Valdivia et al. \(2014\)](#).

We included each of the predictor variables listed in [Table 1](#) as fixed terms in the model. To account for spatial autocorrelation we included sampling unit, nested within the most common habitat type, as random terms in the model, including an offset to account for the length of the sampling unit. We checked for patterns in plots of fitted values against residuals to validate the model. Although there may be a slight trend in decreasing residual variation at higher fitted values ([Fig. S2](#)), indicating that there could be a more optimal link function, this trend was not excessive and the plot was similar to in other published studies ([Valdivia et al., 2014](#)). We also confirmed that there were no patterns evident in plots of predictor variables against residuals. We determined that spatial autocorrelation was not a problem in our models by performing Mantel tests ([Mantel, 1967](#)) ([Fig. S3](#)) and by inspecting the sample variogram for residuals ([Fig. S4](#)). Overdispersion did not appear to be a problem for the final model ($\theta = 1.08$).




Figure 3 Coefficient estimates showing the effect of predictor variables on serval roadkill counts. Error bars represent 95% confidence intervals. We modelled roadkill counts using a generalized linear mixed effect model with a negative binomial distribution and log link. Coefficients with 95% confidence intervals that overlap zero are shown in blue, and those that do not overlap zero are highlighted in green. The full model summary is provided in [Output S5](#).

Full-size [DOI: 10.7717/peerj.6650/fig-3](https://doi.org/10.7717/peerj.6650/fig-3)

RESULTS

A total of 86 serval roadkill events were recorded along the study route between 01/01/2014 to 31/12/2017 ([Fig. 2](#)), which is equivalent to 5.24 carcasses/100 km/year. Our model showed support for effects of season, wetland, and NDVI on serval roadkill counts ([Fig. 3](#), [Output S1](#)). The number of serval roadkills were greater in the dry season than the wet season, and in areas with more wetland, and with greater NDVI ([Fig. 4](#)). There was no support for the effects of the number of guineaowl roadkills, or the anthropogenic variables traffic volume, speed limit, road width, or the amount of road infrastructure on serval roadkill ([Fig. 3](#), [Output S1](#)).

DISCUSSION

Conducting twice-daily transects along both lanes of the 410 km study route, solely to collect roadkill count data for research purposes would have been prohibitively costly. Partnering with the road agency facilitated efficient collection of the necessary data in a standardised manner ([Collinson et al., 2014](#)). Our findings demonstrate that conservation or wildlife




Figure 4 The relationship between serval roadkill counts in each 10 km sampling unit and (A) season, (B) proportion of wetland; and (C) NDVI on the N3 between 2014 and 2017. Boxes (A) and shaded areas (B, C) show 95% confidence intervals.

Full-size [DOI: 10.7717/peerj.6650/fig-4](https://doi.org/10.7717/peerj.6650/fig-4)

management organisations partnering with road patrol agencies can be a powerful strategy for assessing the various factors associated with roadkill rates. Partnerships could help to efficiently reduce roadkill rates and guide future mitigation measures which would benefit road users and contribute to maintaining biodiversity in a developing landscape (Abra *et al.*, 2018).

We calculated an average of 5.24 serval carcasses/100 km/year (85 carcasses in total) on the N3, but there are few studies with which this can be compared. Serval were among the most common species recorded in a previous study on wildlife roadkill on the N3 (14.8% of 183 citizen scientist records, and 1.5% of 209 road patrol records between 2011 and 2014) (Périquet *et al.*, 2018). Five of 17 collared serval were killed by snaring, hunting dogs, and WVCs in the Drakensberg Midlands, South Africa (Ramesh, Kalle & Downs, 2016), but the breakdown of these mortality sources is unclear. Serval roadkill has also been occasionally recorded elsewhere in Africa, with two serval carcasses recorded in a study on roadkill in Uganda (Cibot *et al.*, 2015), but without data on survey effort this cannot be converted into a rate for comparison.

The rate of serval roadkill on the N3 is roughly in line with roadkill rates of other carnivores such as spotted hyaena (*Crocuta crocuta*), common genet (*Genetta genetta*), and black-backed jackal (*Canis mesomelas*) in the Tarangire–Manyara ecosystem, Tanzania (each at approximately 5 individuals/100km/year) (Kioko *et al.*, 2015). Carnivore roadkill rates in Portugal were also similar (5–6 individuals/100 km/year) for species including Egyptian mongoose (*Herpestes ichneumon*), Eurasian badger (*Meles meles*), and common genet (Grilo, Bissonette & Santos-Reis, 2009). However, carnivore roadkill rates can vary

considerably in relation to factors such as species behaviour, abundance, and road placement, for example from 1 to 20 individuals/100 km/year for Western polecats (*Mustela putorius*) and red fox (*Vulpes vulpes*) respectively ([Grilo, Bissonette & Santos-Reis, 2009](#)).

We found support for the hypothesis that serval roadkill counts would be higher in the dry season than the wet season, as carnivores tend to have larger home ranges when water is scarce ([Macdonald & Loveridge, 2010](#)), which could put them at greater risk of WVCs. We also found support for the hypothesis that serval roadkill counts would be positively related to the amount of wetland, which suggests that servals are being killed on roads located in their preferred habitat ([Ramesh & Downs, 2015b](#)), probably because they are more abundant in these areas ([D'Amico et al., 2015](#)). Similar findings were found with the raccoon dog (*Nyctereutes procyonoides viverrinus*) in Japan ([Saeki & Macdonald, 2004](#)), spotted turtles (*Clemmys guttata*) and Blanding's turtles (*Emydoidea blandingii*) in North America ([Beaudry, deMaynadier & Hunter, 2008](#)), and owls in Portugal ([Gomes et al., 2009](#)). Finally, our findings support the hypothesis that there is a positive association between NDVI and roadkill counts of serval. Areas with greater NDVI tend to have higher levels of primary productivity ([Tucker et al., 1985](#)).

There may be a weak relationship between serval roadkill counts and some predictor variables such as infrastructure, and with further data collection this may become more evident. Nevertheless using the current dataset our model did not show support for our other hypotheses, including that serval roadkill counts would be associated with guineafowl roadkill counts, or with anthropogenic variables such as traffic volume, speed limit, road width, or the amount of road infrastructure. This suggests that serval roadkill counts are influenced primarily by ecological drivers such as season, wetland, and NDVI, and that this will vary between species with different ecological requirements. As a result roadkill mitigation strategies may need to be tailored to target species rather than relying on a more generalised approach.

Without data on serval population sizes or other sources of mortality, it is difficult to determine how important WVCs may be for serval population dynamics along the study route. Our findings nevertheless appear to support the suggestion by [Ramesh et al. \(2016\)](#) that WVCs could be a major source of mortality for the species, which is also true for other carnivores. Over 40% of Eurasian badgers in southwest England, for example, were thought to be killed each year in WVCs ([Clarke, White & Harris, 1998](#)). WVCs accounted for 35% of annual mortality of Florida panthers (*Puma concolor coryi*) in the USA ([Taylor et al., 2002](#)), 4–33% of annual mortality of jaguars (*Panthera onca*) in the Atlantic Forest, Brazil ([Cullen Jr et al., 2016](#)), and 17% of annual mortality of Iberian lynx (*Felis pardina*) in southern Spain ([Ferrerias et al., 1992](#)). Identifying factors associated with high roadkill rates is a useful first step in developing roadkill mitigation strategies that could contribute significantly to conservation efforts of some species. Findings from this study suggest that serval roadkills could be reduced if mitigation efforts are focussed on wetland areas and immediate surrounds.

Installation of fencing in combination with wildlife crossing structures ([Clevenger, Chruszcz & Gunson, 2001](#)) like under- and over-passes have been used extensively in

Europe and America ([Danielson & Hubbard, 1998](#); [Goosem et al., 2008](#); [Bissonette & Rosa, 2012](#)), and could be applied to a South African context provided the tendency of serval to use these structures is adequately assessed before implementation. Wildlife crossing structures are widely used by many species ([Clevenger, Chruszcz & Gunson, 2001](#)) including carnivores ([Grilo, Bissonette & Santos-Reis, 2008](#); [Andis, Huijser & Broberg, 2017](#)), and it is reasonable to infer that serval would make use of them as well. The movement patterns of the target species need to be clearly understood, as these are often associated with drainage lines, topography, and habitat ([Clevenger, Chruszcz & Gunson, 2001](#); [Rytwinski et al., 2016](#)). Correct tunnel design at sites optimal for the target species need to be well thought out, as does the installation of the tunnels ([Danielson & Hubbard, 1998](#)). The use of fences must also be carefully considered prior to their use. Fences that are too short in length may exacerbate the problem ([Seiler, 2005](#)) by causing wildlife to follow the fence until the end is reached and there is a gap to cross, thus creating a fence-end hotspot ([Seiler, 2005](#)). Many animals will also either dig under, push through, or jump over fences and consequently collide with vehicles ([Van Niekerk & Eloff, 2005](#)). Caution should be used in applying this approach, because while fencing can be effective at mitigating WVCs, it can also disrupt migration and dispersal routes, reduce gene flow, and exacerbate ecosystem fragmentation ([Creel et al., 2013](#)), while providing materials that can be used by poachers to construct snares ([Lindsey et al., 2011](#); [Williams et al., 2016](#)). The benefits of using of fencing must therefore be carefully weighed against the costs.

Previous research suggests that using the above approach in combination with other methods such as implementing low speed zones, installing traffic-calming devices such as speed bumps ([Glista, DeVault & DeWoody, 2009](#)) or using species-specific warning signage placed in strategic locations to alert road users ([Hardy, Lee & Al-Kaisy, 2006](#)) may increase the effectiveness of mitigation measures ([Grace, Smith & Noss, 2015](#); [Collinson, Marneweck & Davies-Mostert, 2019](#)). While the standard static warning signage typically used is often largely ineffective since drivers quickly habituate to it and fail to make adequate reductions in speed ([Huijser et al., 2015](#)), recent research has shown that when optimised warning signs can be successful at changing driver behaviour and reducing WVCs ([Collinson, Marneweck & Davies-Mostert, 2019](#)). Enhanced warning signs that are specific in time and space may also be effective ([Huijser et al., 2015](#)). While it is possible to modify driver behaviour, our data suggest that the anthropogenic variables included in our models may not be as significant as the various environmental variables in affecting serval roadkill rates on the N3. Removal of roadside vegetation that supports abundant rodent species ([Ruiz-Capillas, Mata & Malo, 2015](#)) could reduce the rate of serval roadkill, although this could impact plant and small mammal conservation. Keeping grass verges trimmed rather than removing vegetation entirely may be effective in manipulating the landscape of fear ([Jacob & Brown, 2000](#)) causing wildlife to spend less time in verge habitats without altering their abundance. If grain spillage from agricultural or transport vehicles is a common occurrence ([Ansara, 2008](#); [Gangadharan et al., 2017](#)), the increased food availability could boost rodent abundance, drawing predators into the road and leading to increased WVCs ([Gangadharan et al., 2017](#)), and this is therefore another factor that needs to be investigated before developing mitigation strategies.

Although useful, there were some caveats to our approach. For example, road patrol teams did not collect data that would have facilitated modelling the number of missed detections (Shilling, Perkins & Collinson, 2015). Furthermore, recording data on which patrol teams conducted which patrols would have enabled us to control for inter-observer bias (Shilling, Perkins & Collinson, 2015), and is recommended for future studies. We also created a WhatsApp group to help observers receive rapid responses to future queries about species identification or any other questions, which we hope will help ensure accurate data collection. We were unable to record the age-sex class of carcasses, which would have been interesting to include in the models (Pallares *et al.*, 2015). We recommend collection of tissue samples to allow this in future studies. Additional data such as the density of serval and prey species at different sampling points along the length of the road would have been incredibly useful in determining the impact on local populations. Finally, the speed at which patrols were driven could bias roadkill detection towards larger species (Collinson *et al.*, 2014), so this method would not be suitable to collect roadkill counts for all species. Further studies incorporating missed carcass detections, inter-observer bias, and population density of target species would be worthwhile. Despite these limitations, this study demonstrated that even the collection of very limited data such as the location, date, and species (or taxon) of roadkill carcasses can help to inform wildlife management policy while minimising additional workload burden on road patrol staff.

CONCLUSIONS

Integrating roadkill data collection into existing road patrols can provide an efficient means of collecting data to allow identification of factors associated with carnivore roadkill. Our findings support the hypotheses that serval roadkill counts were higher in the dry season than the wet season, and they were also higher in areas with more wetland and in areas with greater NDVI. Anthropogenic factors such as traffic volume, speed limit, and the amount of road infrastructure did not influence serval roadkill counts. We suggest that efforts to mitigate serval roadkill, such as installing wildlife crossing structures in combination with fencing, should be targeted at wetlands, but this must be tailored to the ecological requirements of target species.

ACKNOWLEDGEMENTS

We are extremely grateful to N3TC staff for collecting the field data and providing additional data on traffic volumes and speed limits.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Samual T. Williams was supported by a postdoctoral grant from the University of Venda. Lourens H. Swanepoel was supported by the National Research Foundation of South Africa (Grant Nr: 107099 and 115040) and the University of Venda (SMNS/17/Zoo/01). N3 Toll Concession (N3TC) provided financial and logistical support for the research. The funders

had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

University of Venda.

National Research Foundation of South Africa: 107099, 115040.

University of Venda: SMNS/17/Zoo/01.

N3 Toll Concession (N3TC).

Competing Interests

Samual Williams is a Researcher at the Institute for Globally Distributed Open Research and Education (IGDORE). Wendy Collinson and Claire Patterson-Abrolat are currently employed by the Endangered Wildlife Trust (EWT). David Marneweck was employed by the EWT during the data collection period and while the manuscript was prepared, but he is not currently employed by the EWT. Lourens Swanepoel is a director at the African Institute for Conservation Ecology and Genetics

Author Contributions

- Samuel T. Williams conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Wendy Collinson, Claire Patterson-Abrolat and David G. Marneweck conceived and designed the experiments, performed the experiments, contributed reagents/materials/-analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Lourens H. Swanepoel conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The dataset is available at Figshare: Williams, Sam; Collinson, Wendy; Patterson-Abrolat, Claire; Marneweck, David G.; H. Swanepoel, Lourens (2019): Serval roadkill dataset. Figshare. <https://doi.org/10.6084/m9.figshare.6833378.v1>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6650#supplemental-information>.

REFERENCES

- Abra FD, Huijser MP, Pereira CS, Ferraz KM. 2018.** How reliable are your data? Verifying species identification of road-killed mammals recorded by road maintenance personnel in São Paulo State, Brazil. *Biological Conservation* **225**:42–52 DOI [10.1016/j.biocon.2018.06.019](https://doi.org/10.1016/j.biocon.2018.06.019).

- Andis AZ, Huijser MP, Broberg L. 2017.** Performance of arch-style road crossing structures from relative movement rates of large mammals. *Frontiers in Ecology and Evolution* 5:122 DOI 10.3389/fevo.2017.00122.
- Ansara T. 2008.** Determining the ecological status and possible anthropogenic impacts on the grass owl (*Tyto capensis*) population in the East Rand Highveld, Gauteng. MSc thesis, University of Johannesburg.
- Bager A, Rosa CA da. 2010.** Priority ranking of road sites for mitigating wildlife roadkill. *Biota Neotropica* 10:149–153 DOI 10.1590/S1676-06032010000400020.
- Bates D, Mächler M, Bolker B, Walker S. 2015.** Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67:1–48 DOI 10.18637/jss.v067.i01.
- Beaudry F, deMaynadier PG, Hunter ML. 2008.** Identifying road mortality threat at multiple spatial scales for semi-aquatic turtles. *Biological Conservation* 141:2550–2563 DOI 10.1016/j.biocon.2008.07.016.
- Benítez-López A, Alkemade R, Verweij PA. 2010.** The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biological Conservation* 143:1307–1316 DOI 10.1016/j.biocon.2010.02.009.
- Bissonette JA, Rosa S. 2012.** An evaluation of a mitigation strategy for deer-vehicle collisions. *Wildlife Biology* 18:414–423 DOI 10.2981/11-122.
- Bowland JM. 1990.** Diet, home range and movement patterns of serval on farmland in Natal. MSc thesis, University of KwaZulu Natal.
- Cibot M, Bortolamiol S, Seguya A, Krief S. 2015.** Chimpanzees facing a dangerous situation: a high-traffic asphalted road in the Sebitoli area of Kibale National Park, Uganda. *American Journal of Primatology* 77:890–900 DOI 10.1002/ajp.22417.
- Clarke GP, White PCL, Harris S. 1998.** Effects of roads on badger *Meles meles* populations in south-west England. *Biological Conservation* 86:117–124 DOI 10.1016/S0006-3207(98)00018-4.
- Clevenger AP, Chruszcz B, Gunson KE. 2001.** Highway mitigation fencing reduces wildlife-vehicle collisions. *Wildlife Society Bulletin* 29:646–653.
- Collinson WJ, Marneweck C, Davies-Mostert HT. 2019.** Protecting the protected: reducing wildlife roadkill in protected areas. *Animal Conservation* Epub ahead of print Jan 30 2019 DOI 10.1111/acv.12481.
- Collinson WJ, Parker DM, Bernard RTF, Reilly BK, Davies-Mostert HT. 2014.** Wildlife road traffic accidents: a standardized protocol for counting flattened fauna. *Ecology and Evolution* 4:3060–3071 DOI 10.1002/ece3.1097.
- Collinson W, Parker D, Patterson-Abrolat C, Alexander G, Davies-Mostert H. 2015a.** Setjhaba SA, South Afrika: a South African perspective on an emerging transport infrastructure. In: Van der Ree R, Smith DJ, Grilo C, eds. *Handbook of road ecology*. Chichester: John Wiley & Sons, 439–447.
- Collinson WJ, Reilly BK, Parker DM, Bernard RTF, Davies-Mostert HT. 2015b.** An inventory of vertebrate roadkill in the greater Mapungubwe Transfrontier conservation area, South Africa. *South African Journal of Wildlife Research* 45:301–311 DOI 10.3957/056.045.0301.

- Conrad CC, Hilchey KG. 2011. A review of citizen science and community-based environmental monitoring: issues and opportunities. *Environmental Monitoring and Assessment* 176:273–291 DOI 10.1007/s10661-010-1582-5.
- Cook C, Reason CJC, Hewitson BC. 2004. Wet and dry spells within particularly wet and dry summers in the South African summer rainfall region. *Climate Research* 26:17–31 DOI 10.3354/cr026017.
- Creel S, Becker MS, Durant SM, M’Soka J, Matandiko W, Dickman AJ, Christianson D, Dröge E, Mweetwa T, Pettorelli N, Rosenblatt E, Schuette P, Woodroffe R, Bashir S, Beudels-Jamar RC, Blake S, Borner M, Breitenmoser C, Broekhuis F, Cozzi G, Davenport TRB, Deutsch J, Dollar L, Dolrenry S, Douglas-Hamilton I, Fitzherbert E, Foley C, Hazzah L, Henschel P, Hilborn R, Hopcraft JGC, Ikanda D, Jacobson A, Joubert B, Joubert D, Kelly MS, Lichtenfeld L, Mace GM, Milanzi J, Mitchell N, Msuha M, Muir R, Nyahongo J, Pimm S, Purchase G, Schenck C, Sillero-Zubiri C, Sinclair ARE, Songorwa AN, Stanley-Price M, Tehou CA, Trout C, Wall J, Wittemyer G, Zimmermann A. 2013. Conserving large populations of lions—the argument for fences has holes. *Ecology Letters* 16:1413 DOI 10.1111/ele.12145.
- Cullen Jr L, Stanton JC, Lima F, Uezu A, Perilli MLL, Akçakaya HR. 2016. Implications of fine-grained habitat fragmentation and road mortality for jaguar conservation in the Atlantic Forest, Brazil. *PLOS ONE* 11:e0167372 DOI 10.1371/journal.pone.0167372.
- D’Amico M, Román J, De los Reyes L, Revilla E. 2015. Vertebrate road-kill patterns in Mediterranean habitats: who, when and where. *Biological Conservation* 191:234–242 DOI 10.1016/j.biocon.2015.06.010.
- Danielson BJ, Hubbard MW. 1998. A literature review for assessing the status of current methods of reducing deer-vehicle collisions. Iowa Department of Transportation, and the Iowa Department of Natural Resources.
- Didan K. 2015. MOD13Q1 MODIS/Terra Vegetation Indices 16-Day L3 Global 250m SIN Grid V006 [Data set]. Available at https://lpdaac.usgs.gov/dataset_discovery/modis/modis_products_table/mod13q1_v006.
- European Space Agency. 2017. Satellite imagery. Image date 01 December 2017, Available at <https://sentinel.esa.int/web/sentinel/sentinel-data-access>.
- Ferreras P, Aldama JJ, Beltrán JF, Delibes M. 1992. Rates and causes of mortality in a fragmented population of Iberian lynx *Felis pardina* Temminck, 1824. *Biological Conservation* 61:197–202 DOI 10.1016/0006-3207(92)91116-A.
- Forman RTT, Sperling D, Bissonette JA, Clevenger AP, Cutshall CD, Dale VH, Fahrig L, France RL, Heanue K, Goldman CR, Jones J, Swanson F, Turrentine T, Winter TC. 2003. *Road ecology: science and solutions*. Washington, D.C.: Island Press.
- Gangadharan A, Pollock S, Gilhooly P, Friesen A, Dorsey B, St Clair CC. 2017. Grain spilled from moving trains create a substantial wildlife attractant in protected areas. *Animal Conservation* 20:391–400 DOI 10.1111/acv.12336.
- Geertsema AA. 1985. Aspects of the ecology of the serval *Leptailurus serval* in the Ngorongoro Crater, Tanzania. *Netherlands Journal of Zoology* 35:527–610 DOI 10.1163/002829685X00217.

- Geoterraimage. 2015.** 2013–2014 South African National Land-Cover Dataset version 5. Available at http://bgis.sanbi.org/DEA_Landcover/project.asp.
- Glista DJ, DeVault TL, DeWoody JA. 2009.** A review of mitigation measures for reducing wildlife mortality on roadways. *Landscape and Urban Planning* **91**:1–7 DOI [10.1016/j.landurbplan.2008.11.001](https://doi.org/10.1016/j.landurbplan.2008.11.001).
- Gomes L, Grilo C, Silva C, Mira A. 2009.** Identification methods and deterministic factors of owl roadkill hotspot locations in Mediterranean landscapes. *Ecological Research* **24**:355–370 DOI [10.1007/s11284-008-0515-z](https://doi.org/10.1007/s11284-008-0515-z).
- Goosem M, Izumi Y, Turton S. 2001.** Efforts to restore habitat connectivity for an upland tropical rainforest fauna: a trial of underpasses below roads. *Ecological Management & Restoration* **2**:196–202 DOI [10.1046/j.1442-8903.2001.00084.x](https://doi.org/10.1046/j.1442-8903.2001.00084.x).
- Goosem M, Wilson R, Weston N, Cohen M. 2008.** *Highway overpass evaluation of effectiveness: Kuranda Range road upgrade project*. Cairns: James Cook University.
- Grace MK, Smith DJ, Noss RF. 2015.** Testing alternative designs for a roadside animal detection system using a driving simulator. *Nature Conservation* **11**:61–77 DOI [10.3897/natureconservation.11.4420](https://doi.org/10.3897/natureconservation.11.4420).
- Greenwood JD. 2012.** Citizens, science, and environmental policy: a british perspective. In: Dickinson JL, Bonney R, eds. *Citizen science: public participation in environmental research*. Ithaca: Cornell University Press, 150–164.
- Gregory RD, Van Strien A. 2010.** Wild bird indicators: using composite population trends of birds as measures of environmental health. *Ornithological Science* **9**:3–22 DOI [10.2326/osj.9.3](https://doi.org/10.2326/osj.9.3).
- Grilo C, Bissonette JA, Santos-Reis M. 2008.** Response of carnivores to existing highway culverts and underpasses: implications for road planning and mitigation. *Biodiversity and Conservation* **17**:1685–1699 DOI [10.1007/s10531-008-9374-8](https://doi.org/10.1007/s10531-008-9374-8).
- Grilo C, Bissonette JA, Santos-Reis M. 2009.** Spatial-temporal patterns in Mediterranean carnivore road casualties: consequences for mitigation. *Biological Conservation* **142**:301–313 DOI [10.1016/j.biocon.2008.10.026](https://doi.org/10.1016/j.biocon.2008.10.026).
- Guinard É, Prodon R, Barbraud C. 2015.** Case study: a robust method to obtain defendable data on wildlife mortality. In: Van der Ree R, Smith DJ, Grilo C, eds. *Handbook of road ecology*. Chichester: John Wiley & Sons, 96–100.
- Gunson K, Teixeira FZ. 2015.** Road—wildlife mitigation planning can be improved by identifying the patterns and processes associated with wildlife-vehicle collisions. In: Van der Ree R, Smith DJ, Grilo C, eds. *Handbook of road ecology*. Chichester: John Wiley & Sons, 101–109.
- Hardy A, Lee S, Al-Kaisy A. 2006.** Effectiveness of animal advisory messages on dynamic message signs as a speed reduction tool: case study in rural Montana. *Transportation Research Record* **1973**:64–72 DOI [10.3141/1973-10](https://doi.org/10.3141/1973-10).
- Huijser M, Mosler-Berger C, Olsson M, Strein M. 2015.** Wildlife warning signs and animal detection systems aimed at reducing wildlife-vehicle collisions. In: Van der Ree R, Smith DJ, Grilo C, eds. *Handbook of road ecology*. Chichester: John Wiley & Sons, 198–212.

- Jackson SD, Griffin CR. 2000.** A strategy for mitigating highway impacts on wildlife. In: Messmer TA, West B, eds. *Wildlife and highways: seeking solutions to an ecological and socio-economic dilemma*. Nashville: The Wildlife Society, 143–159.
- Jacob J, Brown JS. 2000.** Microhabitat use, giving-up densities and temporal activity as short-and long-term anti-predator behaviors in common voles. *Oikos* **91**:131–138 DOI [10.1034/j.1600-0706.2000.910112.x](https://doi.org/10.1034/j.1600-0706.2000.910112.x).
- Karani P. 2008.** *Impacts of roads on the environment in South Africa*. Midrand: Development Bank of Southern Africa.
- Keshkamat SS, Looijen JM, Zuidgeest MHP. 2009.** The formulation and evaluation of transport route planning alternatives: a spatial decision support system for the Via Baltica project, Poland. *Journal of Transport Geography* **17**:54–64 DOI [10.1016/j.jtrangeo.2008.04.010](https://doi.org/10.1016/j.jtrangeo.2008.04.010).
- Kioko J, Kiffner C, Jenkins N, Collinson WJ. 2015.** Wildlife roadkill patterns on a major highway in northern Tanzania. *African Zoology* **50**:17–22 DOI [10.1080/15627020.2015.1021161](https://doi.org/10.1080/15627020.2015.1021161).
- Laurance WF, Sloan S, Weng L, Sayer JA. 2015.** Estimating the environmental costs of Africa’s massive “development corridors”. *Current Biology* **25**:3202–3208 DOI [10.1016/j.cub.2015.10.046](https://doi.org/10.1016/j.cub.2015.10.046).
- Liang J, Liu Y, Ying L, Li P, Xu Y, Shen Z. 2014.** Road impacts on spatial patterns of land use and landscape fragmentation in three parallel rivers region, Yunnan Province, China. *Chinese Geographical Science* **24**:15–27 DOI [10.1007/s11769-014-0652-y](https://doi.org/10.1007/s11769-014-0652-y).
- Lindsey PA, Romañach SS, Tambling CJ, Chartier K, Groom R. 2011.** Ecological and financial impacts of illegal bushmeat trade in Zimbabwe. *Oryx* **45**:96–111 DOI [10.1017/S0030605310000153](https://doi.org/10.1017/S0030605310000153).
- Loock DJE, Williams ST, Emslie KW, Matthews WS, Swanepoel LH. 2018.** High carnivore population density highlights the conservation value of industrialised sites. *Scientific Reports* **8**:16575 DOI [10.1101/260729](https://doi.org/10.1101/260729).
- Macdonald D, Loveridge A. 2010.** *The biology and conservation of wild felids*. Oxford: Oxford University Press.
- Mantel N. 1967.** The detection of disease clustering and a generalized regression approach. *Cancer Research* **27**:209–220.
- Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C. 2011.** Global biodiversity conservation: the critical role of hotspots. In: Zachos FE, Habel JC, eds. *Biodiversity hotspots*. Berlin: Springer, 3–22 DOI [10.1007/978-3-642-20992-5_1](https://doi.org/10.1007/978-3-642-20992-5_1).
- Mucina L, Rutherford MC. 2006.** *The vegetation of South Africa, Lesotho and Swaziland*. Pretoria: South African National Biodiversity Institute.
- N3 Toll Concession. 2018.** About us. Available at <http://www.n3tc.co.za> (accessed on 01 August 2018).
- OpenStreetMap contributors. 2017.** South Africa OpenStreetMap data. Available at <https://download.geofabrik.de/africa/south-africa.html> (accessed on 25 May 2017).
- Pallares E, Manterola C, Conde DA, Colchero F. 2015.** Case study: roads and jaguars in the Mayan forests. In: Van der Ree R, Smith DJ, Grilo C, eds. *Handbook of road ecology*. Chichester: John Wiley & Sons, 313–316.

- Parchizadeh J, Shilling F, Gatta M, Bencini R, Qashqaei AT, Adibi MA, Williams ST. 2018. Roads threaten Asiatic cheetahs in Iran. *Current Biology* 28:R1141–R1142 DOI 10.1016/j.cub.2018.09.005.
- Périquet S, Roxburgh L, Le Roux A, Collinson WJ. 2018. Testing the value of citizen science for roadkill studies: a case study from South Africa. *Frontiers in Ecology and Evolution* 6:1–15 DOI 10.3389/fevo.2018.00015.
- Perz SG, Qiu Y, Xia Y, Southworth J, Sun J, Marsik M, Rocha K, Passos V, Rojas D, Alarcón G, Barnes G, Baraloto C. 2013. Trans-boundary infrastructure and land cover change: highway paving and community-level deforestation in a tri-national frontier in the Amazon. *Land Use Policy* 34:27–41 DOI 10.1016/j.landusepol.2013.01.009.
- QGIS Development Team. 2018. QGIS geographic information System. Open Source Geospatial Foundation project. Available at <http://qgis.org>.
- R Development Core Team. 2018. R: a language and environment for statistical computing. Version 3.5.0. Vienna: R Foundation for statistical computing. Available at <https://www.R-project.org/>.
- Ramesh T, Downs CT. 2013. Impact of farmland use on population density and activity patterns of serval in South Africa. *Journal of Mammalogy* 94:1460–1470 DOI 10.1644/13-MAMM-A-063.1.
- Ramesh T, Downs CT. 2015a. Impact of land use on occupancy and abundance of terrestrial mammals in the Drakensberg Midlands, South Africa. *Journal for Nature Conservation* 23:9–18 DOI 10.1016/j.jnc.2014.12.001.
- Ramesh T, Downs CT. 2015b. Diet of serval (*Leptailurus serval*) on farmlands in the Drakensberg Midlands, South Africa. *Mammalia* 79:399–407 DOI 10.1515/mammalia-2014-0053.
- Ramesh T, Downs CT, Power RJ, Laurence S, Matthews W, Child MF. 2016. A conservation assessment of *Leptailurus serval*. In: Do Linh San E, Raimondo D, Davies-Mostert HT CMRL, eds. *The Red List of Mammals of South Africa, Swaziland and Lesotho*. Johannesburg: South African National Biodiversity Institute and Endangered Wildlife Trust.
- Ramesh T, Kalle R, Downs CT. 2015. Sex-specific indicators of landscape use by servals: consequences of living in fragmented landscapes. *Ecological Indicators* 52:8–15 DOI 10.1016/j.ecolind.2014.11.021.
- Ramesh T, Kalle R, Downs CT. 2016. Spatiotemporal variation in resource selection of servals: insights from a landscape under heavy land-use transformation. *Journal of Mammalogy* 97:554–567 DOI 10.1093/jmammal/gyv201.
- Ruiz-Capillas P, Mata C, Malo JE. 2015. How many rodents die on the road? Biological and methodological implications from a small mammals' roadkill assessment on a Spanish motorway. *Ecological Research* 30:417–427 DOI 10.1007/s11284-014-1235-1.
- Rytwinski T, Van der Ree R, Cunningham GM, Fahrig L, Findlay CS, Houlihan J, Jaeger JAG, Soanes K, Van der Grift EA. 2015. Experimental study designs to improve the evaluation of road mitigation measures for wildlife. *Journal of Environmental Management* 154:48–64 DOI 10.1016/j.jenvman.2015.01.048.

- Rytwinski T, Soanes K, Jaeger JAG, Fahrig L, Findlay CS, Houlahan J, Van Der Ree R, Van der Grift EA. 2016. How effective is road mitigation at reducing road-kill? A meta-analysis. *PLOS ONE* 11:e0166941 DOI 10.1371/journal.pone.0166941.
- Saeki M, Macdonald DW. 2004. The effects of traffic on the raccoon dog (*Nyctereutes procyonoides viverrinus*) and other mammals in Japan. *Biological Conservation* 118:559–571 DOI 10.1016/j.biocon.2003.10.004.
- Seiler A. 2005. Predicting locations of moose–vehicle collisions in Sweden. *Journal of Applied Ecology* 42(2):371–382 DOI 10.1111/j.1365-2664.2005.01013.x.
- Shilling F, Perkins SE, Collinson W. 2015. Wildlife/roadkill observation and reporting systems. In: Van der Ree R, Smith DJ, Grilo C, eds. *Handbook of road ecology*. Chichester: John Wiley & Sons, 492–501.
- Skinner JD, Chimimba CT (eds.) 2005. *The mammals of the Southern African sub-region*. Cambridge: Cambridge University Press DOI 10.1017/cbo9781107340992.
- Taylor SK, Buergelt CD, Roelke-Parker ME, Homer BL, Rotstein DS. 2002. Causes of mortality of free-ranging Florida panthers. *Journal of Wildlife Diseases* 38:107–114 DOI 10.7589/0090-3558-38.1.107.
- Thiel C. 2011. Ecology and population status of the serval *Leptailurus serval* (Schreber, 1776) in Zambia. PhD Thesis, University of Bonn.
- Thiel C. 2015. *Leptailurus serval*. UCN Red List of Threatened Species. Available at <https://www.iucnredlist.org/species/11638/50654625> (accessed on 20 July 2018).
- Tilman D, Clark M, Williams DR, Kimmel K, Polasky S, Packer C. 2017. Future threats to biodiversity and pathways to their prevention. *Nature* 546:73–81 DOI 10.1038/nature22900.
- Tucker CJ, Vanpraet CL, Sharman MJ, Van Ittersum G. 1985. Satellite remote sensing of total herbaceous biomass production in the Senegalese Sahel: 1980–1984. *Remote Sensing of Environment* 17:233–249 DOI 10.1016/0034-4257(85)90097-5.
- United Nations. 2015. World population prospects: the 2015 revision. Working Paper No. ESA/P/WP241. New York: United Nations. Available at <http://www.un.org/en/development/desa/publications/world-population-prospects-2015-revision.html>.
- Valdivia A, Bruno JF, Cox CE, Hackerott S, Green SJ. 2014. Re-examining the relationship between invasive lionfish and native grouper in the Caribbean. *PeerJ* 2:e348 DOI 10.7717/peerj.348.
- Van Niekerk A, Eloff PJ. 2005. Game, fences and motor vehicle accidents: spatial patterns in the Eastern Cape. *South African Journal of Wildlife Research* 35(2):125–130.
- Verster T, Fourie E. 2018. The good, the bad and the ugly of South African fatal road accidents. *South African Journal of Science* 114:1–7 DOI 10.17159/sajs.2018/20170427.
- Williams S, Collinson W, Patterson-Abrolat C, Marneweck DG, Swanepoel HL. 2018. Serval roadkill dataset. Figshare. DOI 10.6084/m9.figshare.6833378.
- Williams ST, Williams KS, Joubert CJ, Hill RA. 2016. The impact of land reform on the status of large carnivores in Zimbabwe. *PeerJ* 4:e1537 DOI 10.7717/peerj.1537.